# FROM A REMARKABLE MANIFESTATION OF POLAR EFFECTS IN A RADICAL FRAGMENTATION TO THE SYNTHESIS OF HIGHLY FUNCTIONALIZED KETONES




#### Laurent DEBIEN,\* Samir Z. Zard\*


Laboratoire de Synthèse Organique (DCSO), UMR CNRS 7652 École Polytechnique, 91128 Palaiseau Cedex – France




### Precedent Work: Radical Allylation of Simple Allylic Alcohols


- ☐ Process featuring:
  - The **formation** of a C–C  $\sigma$ -bond and a C=C  $\pi$ -bond
  - The **homolytic cleavage** of the usually strong C–O bond














## Scope of the Radical Allylation of Xanthates: Selected Examples


- Method applied to the synthesis of (E)-vinylsulfones, stereodefined alkenes, enol ethers and thioethers, alkylidene cyclubutanes, fluoro- and trifluoromethyl-olefins
- Exceptional functional group tolerance and operational simplicity
- Over 100 functionalized olefins prepared using this method to date!














OL 2011, 13, 5676.




Unpublished result



OL 2011, 13, 6296.




74% yielc  $E/Z = 2:8, dr \ge 95:5$ *OL* **2012**, *14*, 5118.



69% yield E/Z = 8:2*OL* **2012**, *14*, 5118.

# Discovery of a New Mode of Fragmentation: C-O versus C-C Cleavage




□ Polar R groups trigger a rare C–C bond fragmentation

Free alcohol

☐ Identification of the cumyl radical as a **new radical leaving group** 


#### New Radical Allylation for the Synthesis of 1,5-Diketones


Newly formed C-C bond




# **Preparation of Complex Structures**

Further demonstration of the process tolerance









# Acknowledgements

- Pr. Samir S. Zard
- ☐ Zard group
- ☐ Funding and organizing agencies







